

LIE ALGEBRAS ASSOCIATED TO HYPERPLANE ARRANGEMENTS

Alexandru Suciu

Northeastern University

Workshop: Various Guises of Reflection Arrangements

International Centre for Mathematical Sciences, Edinburgh, UK

March 14, 2023

LOWER CENTRAL SERIES

- ▶ Let G be a group. The *lower central series* $\{\gamma_k(G)\}_{k \geq 1}$ is defined inductively by $\gamma_1(G) = G$ and $\gamma_{k+1}(G) = [G, \gamma_k(G)]$.
- ▶ Here, if $H, K < G$, then $[H, K]$ is the subgroup of G generated by $\{[a, b] := aba^{-1}b^{-1} \mid a \in H, b \in K\}$. If $H, K \triangleleft G$, then $[H, K] \triangleleft G$.
- ▶ The subgroups $\gamma_k(G)$ are, in fact, characteristic subgroups of G . Moreover, $[\gamma_k(G), \gamma_\ell(G)] \subseteq \gamma_{k+\ell}(G)$, $\forall k, \ell \geq 1$.
- ▶ In particular, it is a *central* series, i.e., $[G, \gamma_k(G)] \subseteq \gamma_{k+1}(G)$.
- ▶ In fact, it is the fastest descending central series for G .
- ▶ It is also a *normal* series, i.e., $\gamma_k(G) \triangleleft G$. Each quotient,

$$\text{gr}_k(G) := \gamma_k(G) / \gamma_{k+1}(G)$$

lies in the center of $G / \gamma_{k+1}(G)$, and thus is an abelian group.

- ▶ If G is finitely generated, then so are its LCS quotients. Set $\phi_k(G) := \text{rank gr}_k(G)$.

ASSOCIATED GRADED LIE ALGEBRA

- ▶ For a coefficient ring \mathbb{k} , we let $\text{gr}(\mathbf{G}; \mathbb{k}) = \bigoplus_{k \geq 1} \text{gr}_k(\mathbf{G}) \otimes \mathbb{k}$.
- ▶ This is a graded Lie algebra, with addition induced by the group multiplication and with Lie bracket $[\cdot, \cdot]: \text{gr}_k \times \text{gr}_\ell \rightarrow \text{gr}_{k+\ell}$ induced by the group commutator.
- ▶ The construction is functorial. Write $\text{gr}(\mathbf{G}) = \text{gr}(\mathbf{G}; \mathbb{Z})$.
- ▶ Example: if F_n is the free group of rank n , then
 - $\text{gr}(F_n)$ is the free Lie algebra $\text{Lie}(\mathbb{Z}^n)$.
 - $\text{gr}_k(F_n)$ is free abelian, of rank $\phi_k(F_n) = \frac{1}{k} \sum_{d|k} \mu(d) n^{\frac{k}{d}}$.
- ▶ $\mathbf{G}/\gamma_k(\mathbf{G})$ is the maximal $(k-1)$ -step nilpotent quotient of \mathbf{G} .
- ▶ $\mathbf{G}/\gamma_2(\mathbf{G}) = \mathbf{G}_{\text{ab}}$, while $\mathbf{G}/\gamma_3(\mathbf{G}) \leftrightarrow H^{\leq 2}(\mathbf{G}; \mathbb{Z})$.

CHEN LIE ALGEBRAS

- ▶ Let $G^{(i)}$ be the *derived series* of G , starting at $G^{(1)} = G'$, $G^{(2)} = G''$, and defined inductively by $G^{(i+1)} = [G^{(i)}, G^{(i)}]$.
- ▶ The quotient groups, $G/G^{(i)}$, are solvable; $G/G' = G_{ab}$, while G/G'' is the maximal metabelian quotient of G .
- ▶ The i -th *Chen Lie algebra* of G is defined as $\text{gr}(G/G^{(i)}; \mathbb{k})$.
- ▶ The projection $q_i: G \twoheadrightarrow G/G^{(i)}$, induces a surjection $\text{gr}_k(G; \mathbb{k}) \twoheadrightarrow \text{gr}_k(G/G^{(i)}; \mathbb{k})$, which is an iso for $k \leq 2^i - 1$.
- ▶ Assuming G is finitely generated, write $\theta_k(G) = \text{rank } \text{gr}_k(G/G'')$ for the *Chen ranks*. We have $\phi_k(G) \geq \theta_k(G)$, with equality for $k \leq 3$.
- ▶ Example (K.-T. Chen 1951): $\theta_k(F_n) = (k-1) \binom{n+k-2}{k}$, for $k \geq 2$.

HOLONOMY LIE ALGEBRA

- ▶ A quadratic approximation of the Lie algebra $\text{gr}(\mathbf{G}; \mathbb{k})$, where \mathbb{k} is a field, is the *holonomy Lie algebra* of \mathbf{G} , defined as

$$\mathfrak{h}(\mathbf{G}; \mathbb{k}) := \text{Lie}(H_1(\mathbf{G}; \mathbb{k})) / \langle \text{im}(\mu_{\mathbf{G}}^{\vee}) \rangle,$$

where



- $L = \text{Lie}(V)$ the free Lie algebra on the \mathbb{k} -vector space $V = H_1(\mathbf{G}; \mathbb{k})$, with $L_1 = V$ and $L_2 = V \wedge V$;
 - $\mu_{\mathbf{G}}^{\vee}: H_2(\mathbf{G}; \mathbb{k}) \rightarrow V \wedge V$ is the dual of the cup product map $\mu_{\mathbf{G}}: H^1(\mathbf{G}; \mathbb{k}) \wedge H^1(\mathbf{G}; \mathbb{k}) \rightarrow H^2(\mathbf{G}; \mathbb{k})$.
- ▶ There is natural epimorphism of graded Lie algebras, $\mathfrak{h}(\mathbf{G}; \mathbb{k}) \twoheadrightarrow \text{gr}(\mathbf{G}; \mathbb{k})$, which restricts to isos in degrees 1 and 2.
 - ▶ For each $i \geq 2$, this morphism factors through epimorphisms $\mathfrak{h}(\mathbf{G}; \mathbb{k}) / \mathfrak{h}(\mathbf{G}; \mathbb{k})^{(i)} \twoheadrightarrow \text{gr}(\mathbf{G}/\mathbf{G}^{(i)}; \mathbb{k})$.

MALCEV LIE ALGEBRA

- ▶ Let \mathbb{k} be a field of characteristic 0. Then $\mathbb{k}G$ is a Hopf algebra, with comultiplication $\Delta(g) = g \otimes g$ and counit $\varepsilon: \mathbb{k}G \rightarrow \mathbb{k}$.
- ▶ Let $J = \ker \varepsilon$. The J -adic completion $\widehat{\mathbb{k}G} = \varprojlim_k \mathbb{k}G/J^k$ is a filtered, complete Hopf algebra.
- ▶ An element $x \in \widehat{\mathbb{k}G}$ is called *primitive* if $\widehat{\Delta}x = x \widehat{\otimes} 1 + 1 \widehat{\otimes} x$. The set $\mathfrak{m}(G; \mathbb{k}) = \text{Prim}(\widehat{\mathbb{k}G})$ of all such elements, with bracket $[x, y] = xy - yx$, is the *Malcev Lie algebra* of G .
- ▶ If G is finitely generated, then $\text{gr}(\mathfrak{m}(G; \mathbb{k})) \cong \text{gr}(G; \mathbb{k})$.
- ▶ G is *filtered-formal* (over \mathbb{k}), if there is an isomorphism of filtered Lie algebras, $\mathfrak{m}(G; \mathbb{k}) \cong \widehat{\text{gr}}(G; \mathbb{k})$.
- ▶ G is *1-formal* (over \mathbb{k}) if it is filtered formal and the projection $\mathfrak{h}(G; \mathbb{k}) \rightarrow \text{gr}(G; \mathbb{k})$ is an isomorphism; that is, $\mathfrak{m}(G; \mathbb{k}) \cong \widehat{\mathfrak{h}}(G; \mathbb{k})$.
- ▶ (Papadima–S. 2004) If G is 1-formal, then the maps $\mathfrak{h}(G; \mathbb{k})/\mathfrak{h}(G; \mathbb{k})^{(i)} \rightarrow \text{gr}(G/G^{(i)}; \mathbb{k})$ are isomorphisms.

HYPERPLANE ARRANGEMENTS

- ▶ An *arrangement of hyperplanes* is a finite collection \mathcal{A} of codimension 1 linear (or affine) subspaces in \mathbb{C}^d .
- ▶ For each $H \in \mathcal{A}$ let α_H be a linear form with $\ker(\alpha_H) = H$; set $f = \prod_{H \in \mathcal{A}} \alpha_H$.
- ▶ *Intersection lattice* $L(\mathcal{A})$: poset of all intersections of \mathcal{A} , ordered by reverse inclusion, and ranked by codimension.



- ▶ *Complement*: $M(\mathcal{A}) = \mathbb{C}^d \setminus \bigcup_{H \in \mathcal{A}} H$. It is a smooth, quasi-projective variety and also a Stein manifold. It has the homotopy type of a finite, connected, d -dimensional CW-complex.

EXAMPLE (THE BOOLEAN ARRANGEMENT)

- ▶ \mathfrak{B}_n : all coordinate hyperplanes $z_i = 0$ in \mathbb{C}^n .
- ▶ $L(\mathfrak{B}_n)$: Boolean lattice of subsets of $\{0, 1\}^n$.
- ▶ $M(\mathfrak{B}_n)$: complex algebraic torus $(\mathbb{C}^*)^n \simeq K(\mathbb{Z}^n, 1)$.

EXAMPLE (THE BRAID ARRANGEMENT)


- ▶ \mathcal{A}_n : all diagonal hyperplanes $z_i - z_j = 0$ in \mathbb{C}^n .
- ▶ $L(\mathcal{A}_n)$: lattice of partitions of $[n] := \{1, \dots, n\}$, ordered by refinement.
- ▶ $M(\mathcal{A}_n)$: the (ordered) configuration space of n distinct points in \mathbb{C} ; it is a classifying space $K(P_n, 1)$ for the pure braid group on n strands, P_n .

COHOMOLOGY RINGS OF ARRANGEMENTS

- ▶ The space $M(\mathcal{A})$ admits a minimal cell structure.
- ▶ The groups $H_q(M(\mathcal{A}); \mathbb{Z})$ are finitely generated and torsion-free, with ranks given by $\sum_{q=0}^{\ell} b_q(M(\mathcal{A}))t^q = \sum_{X \in L(\mathcal{A})} \mu(X)(-t)^{\text{rank}(X)}$, where $\mu: L(\mathcal{A}) \rightarrow \mathbb{Z}$ is defined by $\mu(\mathbb{C}^d) = 1$ and $\mu(X) = -\sum_{Y \supsetneq X} \mu(Y)$.
- ▶ Let E be the \mathbb{Z} -exterior algebra on degree 1 cohomology classes $e_H = \frac{1}{2\pi i} [d \log(\alpha_H)]$ dual to the meridians x_H around $H \in \mathcal{A}$.
- ▶ Let $\partial: E^* \rightarrow E^{*-1}$ be the differential given by $\partial(e_H) = 1$, and set $e_X = \prod_{H \supsetneq X} e_H$ for each $X \in \mathcal{L}(\mathcal{A})$.
- ▶ Arnold, Brieskorn, Orlik–Solomon showed: $H^*(M(\mathcal{A}); \mathbb{Z}) \cong E/I$, where $I = \langle \partial e_X : \text{rank}(X) < |X| \rangle$.
- ▶ M. Kim and B. Shapiro: The quasi-projective variety M admits a *pure* mixed Hodge structure.
- ▶ Thus, M is \mathbb{Q} -formal (albeit not \mathbb{Z}_p -formal, in general).

FUNDAMENTAL GROUPS OF ARRANGEMENTS

- ▶ Let $\mathcal{A}' = \{H \cap \mathbb{C}^2\}_{H \in \mathcal{A}}$ be a generic planar section of \mathcal{A} . Then the arrangement group, $G(\mathcal{A}) = \pi_1(M(\mathcal{A}))$, is isomorphic to $\pi_1(M(\mathcal{A}'))$.
- ▶ So let \mathcal{A} be an arrangement of n affine lines in \mathbb{C}^2 . Taking a generic projection $\mathbb{C}^2 \rightarrow \mathbb{C}$ yields the braid monodromy $\alpha = (\alpha_1, \dots, \alpha_s)$, where $s = \#\{\text{multiple points}\}$ and the braids $\alpha_r \in P_n$ can be read off an associated braided wiring diagram,



- ▶ The group $G(\mathcal{A})$ has a presentation with meridional generators x_1, \dots, x_n and commutator relators $x_i \alpha_j (x_i)^{-1}$.

HOLONOMY AND ASSOCIATED GRADED LIE ALGEBRAS

- ▶ The holonomy Lie algebra of $G = G(\mathcal{A})$ is determined by $L_{\leq 2}(\mathcal{A})$,

$$\mathfrak{h}(G) = \text{Lie}(x_H : H \in \mathcal{A}) / \text{ideal} \left\{ \left[x_H, \sum_{\substack{K \in \mathcal{A} \\ K \supset Y}} x_K \right] : \begin{array}{l} H \in \mathcal{A}, Y \in L_2(\mathcal{A}) \\ H \supset Y \end{array} \right\}.$$

- ▶ Since M is formal, the group G is 1-formal. Hence, $\text{gr}(G) \otimes \mathbb{Q}$ is determined by $H^{\leq 2}(M, \mathbb{Q})$, and thus, by $L_{\leq 2}(\mathcal{A})$.
- ▶ In fact, the surjection $\mathfrak{h}(G) \twoheadrightarrow \text{gr}(G)$ induces an isomorphism, $\mathfrak{h}(G) \otimes \mathbb{Q} \xrightarrow{\cong} \text{gr}(G) \otimes \mathbb{Q}$.
- ▶ (Papadima–S. 2004) The Chen ranks $\theta_k(G)$ are also determined by $L_{\leq 2}(\mathcal{A})$.
- ▶ Explicit combinatorial formulas for the LCS ranks $\phi_k(G)$ are known in some cases, but not in general.

- ▶ $U(\mathfrak{h}(\mathbf{G}) \otimes \mathbb{Q}) = \text{Ext}_{\mathcal{A}}^1(\mathbb{Q}, \mathbb{Q}) = \overline{\mathcal{A}}^!$, the quadratic dual of the quadratic closure of $\mathcal{A} = H^*(M, \mathbb{Q})$.
- ▶ (Falk–Randell 1985) If \mathcal{A} is *supersolvable* with exponents d_1, \dots, d_ℓ , then $\phi_k(\mathbf{G}) = \sum_{i=1}^{\ell} \phi_k(F_{d_i})$. (Also follows from Koszulity of $H^*(M, \mathbb{Q})$ and Koszul duality.)
- ▶ (Porter–S. 2020) The map $\mathfrak{h}_3(\mathbf{G}) \rightarrow \text{gr}_3(\mathbf{G})$ is an isomorphism, but it is not known whether $\mathfrak{h}_3(\mathbf{G})$ is torsion-free.
- ▶ (S. 2002) The groups $\text{gr}_k(\mathbf{G})$ may have non-zero torsion for $k \gg 0$. E.g., if $\mathbf{G} = \mathbf{G}(\text{MacLane})$, then $\text{gr}_5(\mathbf{G}) = \mathbb{Z}^{87} \oplus \mathbb{Z}_2^4 \oplus \mathbb{Z}_3$.
- ▶ (S. 2002): Is the torsion in $\text{gr}(\mathbf{G})$ combinatorially determined?
- ▶ (Artal Bartolo, Guerville-Ballé, and Viu-Sos 2020): Answer: No!
- ▶ There are two arrangements of 13 lines, \mathcal{A}^\pm , each one with 11 triple points and 2 quintuple points, such that $\text{gr}_k(\mathbf{G}^+) \cong \text{gr}_k(\mathbf{G}^-)$ for $k \leq 3$, yet $\text{gr}_4(\mathbf{G}^+) = \mathbb{Z}^{211} \oplus \mathbb{Z}_2$ and $\text{gr}_4(\mathbf{G}^-) = \mathbb{Z}^{211}$.

NILPOTENT QUOTIENTS

- ▶ The quotient $G/\gamma_3(G)$ is determined by $L_{\leq 2}(\mathcal{A})$. Indeed, in the central extension,

$$0 \longrightarrow \text{gr}_2(G) \longrightarrow G/\gamma_3(G) \longrightarrow G_{\text{ab}} \longrightarrow 0,$$

we have $\text{gr}_2(G) = (I^2)^\vee$ and the k -invariant $H_2(G_{\text{ab}}) \rightarrow \text{gr}_2(G)$ is dual of the inclusion $I^2 \hookrightarrow E^2 = \bigwedge^2 G_{\text{ab}}$.

- ▶ (G. Rybnikov 1994): $G/\gamma_4(G)$ is not always determined by $L_{\leq 2}(\mathcal{A})$.
- ▶ There are two arrangements of 13 lines, \mathcal{A}^\pm , each one with 15 triple points, such that $L(\mathcal{A}^+) \cong L(\mathcal{A}^-)$, and therefore $G^+/\gamma_3(G^+) \cong G^-/\gamma_3(G^-)$ and $\text{gr}_3(G^+) \cong \text{gr}_3(G^-)$, but $G^+/\gamma_4(G^+) \not\cong G^-/\gamma_4(G^-)$.
- ▶ The difference can be explained in terms of (generalized) Massey triple products over \mathbb{Z}_3 .

DECOMPOSABLE ARRANGEMENTS

- ▶ For each flat $X \in L(\mathcal{A})$, let $\mathcal{A}_X := \{H \in \mathcal{A} \mid H \supset X\}$ be the localization of \mathcal{A} at X .
- ▶ The inclusions $\mathcal{A}_X \subset \mathcal{A}$ give rise to maps $M(\mathcal{A}) \hookrightarrow M(\mathcal{A}_X)$. Restricting to rank 2 flats yields a map

$$j: M(\mathcal{A}) \longrightarrow \prod_{X \in L_2(\mathcal{A})} M(\mathcal{A}_X).$$

- ▶ The induced homomorphism on fundamental groups, $j_{\#}$, defines a morphism of graded Lie algebras,

$$\mathfrak{h}(j_{\#}): \mathfrak{h}(\mathcal{G}) \longrightarrow \prod_{X \in L_2(\mathcal{A})} \mathfrak{h}(\mathcal{G}_X).$$

THEOREM (PAPADIMA–S. 2006)

The map $\mathfrak{h}_k(j_{\#})$ is a surjection for each $k \geq 3$ and an iso for $k = 2$.

DEFINITION

\mathcal{A} is *decomposable* if the map $\mathfrak{h}_3(j_{\#})$ is an isomorphism.

EXAMPLE

Let $\mathcal{A}(\Gamma) = \{z_i - z_j = 0 : (i, j) \in E(\Gamma)\} \subset \mathcal{A}_n$ be a graphic arrangement. Then $\mathcal{A}(\Gamma)$ is decomposable if and only if Γ contains no K_4 subgraph.

THEOREM (PAPADIMA–S. 2006)

Let \mathcal{A} be a decomposable arrangement, and let $G = G(\mathcal{A})$. Then

- ▶ The map $\mathfrak{h}'(j_{\#}) : \mathfrak{h}'(G) \rightarrow \prod_{X \in L_2(\mathcal{A})} \mathfrak{h}'(G_X)$ is an isomorphism of graded Lie algebras.
- ▶ The map $\mathfrak{h}(G) \twoheadrightarrow \text{gr}(G)$ is an isomorphism
- ▶ For each $k \geq 2$, the group $\text{gr}_k(G)$ is free abelian of rank $\phi_k(G) = \sum_{X \in L_2(\mathcal{A})} \phi_k(F_{\mu(X)})$.

THEOREM (PORTER–S. 2020)

Let \mathcal{A} and \mathcal{B} be decomposable arrangements with $L_{\leq 2}(\mathcal{A}) \cong L_{\leq 2}(\mathcal{B})$. Then, for each $k \geq 2$,

$$G(\mathcal{A})/\gamma_k(G(\mathcal{A})) \cong G(\mathcal{B})/\gamma_k(G(\mathcal{B})).$$

THE MILNOR FIBRATION



- ▶ The defining polynomial map $f: \mathbb{C}^d \rightarrow \mathbb{C}$ restricts to a smooth fibration, $f: M \rightarrow \mathbb{C}^*$, called the *Milnor fibration* of \mathcal{A} .
- ▶ The *Milnor fiber* is $F(\mathcal{A}) := f^{-1}(1)$. The monodromy, $h: F \rightarrow F$, is given by $h(z) = e^{2\pi i/n}z$, where $n = |\mathcal{A}|$.
- ▶ F is a Stein manifold. It has the homotopy type of a finite CW-complex of dimension $d - 1$ (connected if $d > 1$).
- ▶ (Zuber 2010) MHS on F may not be pure and $\pi_1(F)$ may be non-1-formal.

- ▶ F is the regular, \mathbb{Z}_n -cover of $U = \mathbb{P}(M)$, classified by the epimorphism $\pi_1(U) \twoheadrightarrow \mathbb{Z}_n$, $x_H \mapsto 1$.
- ▶ Let $\iota: F \hookrightarrow M$ be the inclusion. Induced maps on π_1 :

$$\begin{array}{ccccccc}
 & & & & 1 & & \\
 & & & & \downarrow & & \\
 & & & & \mathbb{Z} & & \\
 & & & & \downarrow & & \\
 1 & \longrightarrow & \pi_1(F) & \xrightarrow{\iota_{\#}} & \pi_1(M) & \xrightarrow{f_{\#}} & \mathbb{Z} \longrightarrow 1 \\
 & & & & \downarrow \rho_{\#} & & \\
 & & & & \pi_1(U) & & \\
 & & & & \downarrow & & \\
 & & & & 1 & &
 \end{array}$$

- ▶ $b_1(F) \geq n - 1$, and may be computed from the characteristic varieties $\mathcal{V}_S^1(U)$. Combinatorial formulas are known in some cases, e.g., if $\mathbb{P}(\mathcal{A})$ has only double or triple points (Papadima–S. 2017).
- ▶ (Denham–S. 2016) $H_*(F; \mathbb{Z})$ may have torsion. (Yoshinaga 2020): in fact, $H_1(F; \mathbb{Z})$ may have torsion.

THEOREM (S. 2021)

Suppose $h_*: H_1(F; \mathbb{Z}) \rightarrow H_1(F; \mathbb{Z})$ is the identity. Then


- ▶ $\text{gr}_{\geq 2}(\pi_1(F)) \cong \text{gr}_{\geq 2}(G)$.
- ▶ $\text{gr}_{\geq 2}(\pi_1(F)/\pi_1(F)'') \cong \text{gr}_{\geq 2}(G/G'')$.

THEOREM (S. 2021)

Suppose $h_*: H_1(F, \mathbb{Q}) \rightarrow H_1(F, \mathbb{Q})$ is the identity. Then

- ▶ $\text{gr}_{\geq 2}(\pi_1(F)) \otimes \mathbb{Q} \cong \text{gr}_{\geq 2}(G) \otimes \mathbb{Q}$.
- ▶ $\text{gr}_{\geq 2}(\pi_1(F)/\pi_1(F)'') \otimes \mathbb{Q} \cong \text{gr}_{\geq 2}(G/G'') \otimes \mathbb{Q}$.
- ▶ $\phi_k(\pi_1(F)) = \phi_k(G)$ and $\theta_k(\pi_1(F)) = \theta_k(G)$ for all $k \geq 2$.

FALK'S PAIR OF ARRANGEMENTS






- ▶ Both \mathcal{A} and \mathcal{A}' have 2 triple points and 9 double points, yet $L(\mathcal{A}) \not\cong L(\mathcal{A}')$. Nevertheless, $M(\mathcal{A}) \simeq M(\mathcal{A}')$.
- ▶ Both Milnor fibrations have trivial \mathbb{Z} -monodromy.
- ▶ (S. 2017) $\pi_1(F) \not\cong \pi_1(F')$.
- ▶ The difference is picked by the depth-2 characteristic varieties: $\mathcal{V}_2(F) \cong \mathbb{Z}_3$, yet $\mathcal{V}_2(F') = \{1\}$

YOSHINAGA'S ICOSIDODECAHEDRAL ARRANGEMENT

- ▶ The icosidodecahedron is the convex hull of **30** vertices given by the even permutations of $(0, 0, \pm 1)$ and $\frac{1}{2}(\pm 1, \pm \phi, \pm \phi^2)$, where $\phi = (1 + \sqrt{5})/2$.
- ▶ It gives rise to an arrangement of **16** hyperplanes in \mathbb{R}^3 , whose complexification is the icosidodecahedral arrangement \mathcal{A} in \mathbb{C}^3 .
- ▶ $M(\mathcal{A})$ is a $K(G, 1)$.
- ▶ $H_1(F; \mathbb{Z}) = \mathbb{Z}^{15} \oplus \mathbb{Z}_2$. Thus, the algebraic monodromy of the Milnor fibration is trivial over \mathbb{Q} and \mathbb{Z}_p ($p > 2$), but not over \mathbb{Z} .
- ▶ Hence, $\text{gr}(\pi_1(F)) \cong \text{gr}(\pi_1(U))$, away from the prime **2**. Moreover,
 - $\text{gr}_1(\pi_1(F)) = \mathbb{Z}^{15} \oplus \mathbb{Z}_2$
 - $\text{gr}_2(\pi_1(F)) = \mathbb{Z}^{45} \oplus \mathbb{Z}_2^7$
 - $\text{gr}_3(\pi_1(F)) = \mathbb{Z}^{250} \oplus \mathbb{Z}_2^{43}$
 - $\text{gr}_4(\pi_1(F)) = \mathbb{Z}^{1,405} \oplus \mathbb{Z}_2^7$ and $\mathfrak{h}_4(\pi_1(F)) = \mathbb{Z}^{1,405} \oplus \mathbb{Z}_2^{20}$.

REFERENCES

-  Richard D. Porter and Alexander I. Suci, *Homology, lower central series, and hyperplane arrangements*, *European Journal of Mathematics* **6** (2020), nr. 3, 1039–1072.
-  Alexander I. Suci, *Lower central series and split extensions*, [arXiv:2105.14129](https://arxiv.org/abs/2105.14129).
-  Alexander I. Suci, *Alexander invariants and cohomology jump loci in group extensions*, *Annali della Scuola Normale Superiore di Pisa* (to appear), [doi:10.2422/2036-2145.202112_005](https://doi.org/10.2422/2036-2145.202112_005), [arXiv:2107.05148](https://arxiv.org/abs/2107.05148).